



ADDING JEMPERLI (DOSTARLIMAB-GXLY) TO AN ORACLE HEALTH[®] TREATMENT PLAN

INDICATIONS

- JEMPERLI, in combination with carboplatin and paclitaxel, followed by JEMPERLI as a single agent, is indicated for the treatment of adult patients with primary advanced or recurrent endometrial cancer (EC).
- JEMPERLI, as a single agent, is indicated for the treatment of adult patients with mismatch repair deficient (dMMR) recurrent or advanced:
 - EC, as determined by an FDA-approved test, that has progressed on or following prior treatment with a platinum-containing regimen in any setting and are not candidates for curative surgery or radiation, or
 - solid tumors, as determined by an FDA-approved test, that have progressed on or following prior treatment and who have no satisfactory alternative treatment options. This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

IMPORTANT SAFETY INFORMATION

Severe and Fatal Immune-Mediated Adverse Reactions

- Immune-mediated adverse reactions, which can be severe or fatal, can occur in any organ system or tissue and can occur at any time during or after treatment with a PD-1/PD-L1–blocking antibody, including JEMPERLI.
- Monitor closely for signs and symptoms of immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function tests at baseline and periodically during treatment. For suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.

Please see additional [Important Safety Information](#) on pages 6-8 and full [Prescribing Information](#).

dMMR = mismatch repair deficient.

Oracle Health is a registered trademark of Oracle Corporation.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

ABOUT THIS GUIDE

This electronic medical record (EMR) guide is intended to help health care providers (HCPs) who want to create Treatment Plans that include JEMPERLI (dostarlimab-gxly) or want to add JEMPERLI to an existing Treatment Plan. Treatment Plans group together order sets for medications, lab testing, procedures, and other aspects of care based on the patient's diagnosis and condition. It is important to evaluate oncology treatment plans frequently as treatment options, such as JEMPERLI, become available.

This guide does not constitute guidance for treatment or medical advice. It is the responsibility of the HCP to select a treatment based on their independent medical judgment and the needs of each individual patient.

The examples and instructions listed in this guide are based on the most recent version of Oracle Health. Locations, illustrations, and terminology are subject to change with system updates. This guide is meant to serve as an overview only and should not replace detailed instructions provided to you by your internal or external EMR support resources. GSK makes no claims or warranties about the applicability or appropriateness of this information. This guide has not been reviewed or endorsed by Oracle Health. GSK does not endorse or recommend any EMR system.

EMR TREATMENT PLANS HELP SIMPLIFY ONCOLOGY WORKFLOWS

Treatment Plans are commonly used to help facilitate patient care. After the initial release of a Treatment Plan by a medical society, the Oracle Health system may benefit from a clinical update. The optimization of Treatment Plans is a common process and provides an opportunity to incorporate treatment updates. Treatment Plans are typically modified at the health system level to help reduce treatment variation. Typically, an oncology practice will conduct a clinical review process to confirm and approve a suggested Treatment Plan optimization. Various stakeholders may participate in reviewing Treatment Plan optimization requests prior to the approval.

As treatment options such as JEMPERLI become available, it may be necessary to create a new Treatment Plan or to update an existing Treatment Plan to remove system obstacles to prescribe JEMPERLI for its approved indications. Updating relevant Treatment Plans to include JEMPERLI communicates to the care team that it is available to order for appropriate patients.


NOTE: If JEMPERLI is not available for selection in Oracle Health, the practice may need to run a drug database update. As a backup option, the practice EMR Support/IT Team may be able to manually add JEMPERLI, subject to the practice's business rules for drug database maintenance.

How to Add JEMPERLI to a Treatment Plan

Upon request and approval from the Clinical Team, the Health System IT Team can build Treatment Plans, called PowerPlans, of frequently written groups of orders for easier selection. PowerPlans can be based on published treatment protocols and enable consistency of care and efficiency of ordering. If the practice has existing PowerPlans, it may be efficient to modify an existing PowerPlan to include new therapies. If the practice does not have appropriate existing PowerPlans, a new PowerPlan can be created. Updating existing PowerPlans or adding new PowerPlans is typically managed by the practice EMR Support/IT Team using an established process for requesting, approving, and implementing EMR changes.


Adding Orders to an Existing PowerPlan Phase

- From the DCP Tool, launch the **PowerPlan Tool**.
Select **Open Existing Plan** .




PowerPlan Toolbar

- From the **Plan Selection** window, select the appropriate **Plan**.




Searching for a PowerPlan

- Select **OK**.
- If the PowerPlan has multiple phases, from the **Description** column, select the phase.



Example of a Phase Selection

- Select the **Prescription** tab in the lower-right section of the main window.
- Enter text into the **Start search at** box and click the **Find** button  to search for orderable items, such as JEMPERLI.
- Select the item or items in the **Search results** box to add to the PowerPlan. Click the right arrow  to add the selected orderable item(s) to the **Current list**.



A Medication in the Current list

Adding Orders to an Existing PowerPlan Phase (cont.)

8. Select **Add**.
9. From the **Order Sentence** tab, move all appropriate instructions to the **Current list**.
 - Administration:
For Preparation and Administration, please refer to the [Prescribing Information](#)

Order	Note	Outcome	Order Sentence	Copy Components	Sub Phase	Prescription
Order Sentences: = JEMPERLI injection, for intravenous use = Dose 1 through 4: 500 mg every 3 weeks = Subsequent dosing beginning 3 weeks after Dose 4 (Dose 5 onwards): 1,000 mg every 6 weeks = Administration: For Preparation and Administration, please refer to the Prescribing Information						

Please refer to the [Prescribing Information](#) for the proper dosage of each indication of Jemperli and administration instructions.

Example of Adding Order Sentences to the Medication Order

10. Select **Add** to add items from the **Current list** to the Order Set. The component is displayed in the **Description** column.
11. Select **Save**.
12. Similarly, add other appropriate Orders to complete the Order Set. For example, lab or patient education Orders.
13. Select the phase appropriate for the Order, such as laboratory or patient education.

Order	Note	Outcome	Order Sentence	Copy components	Sub Phase	Prescription						
Start search at: CMP All Facilities Mnemonic type filter: Activity type filter: Search results: Current list: <table border="1"> <thead> <tr> <th>Synonym</th> <th>Clinical Category</th> <th>Clinical Subcategory</th> </tr> </thead> <tbody> <tr> <td></td> <td> </td> <td> </td> </tr> </tbody> </table> Add Reset							Synonym	Clinical Category	Clinical Subcategory			
Synonym	Clinical Category	Clinical Subcategory										

Adding Other Orders to the PowerPlan

14. Select the **Order** tab. Search for and select lab orders.
15. Select **Add** for each additional item, and add Order Sentence as needed.
16. Select **Save** from the toolbar.

DCP: DB PowerPlan Tool

Task Prerequisites Reports Options Help

Description

Order Set Name

- Laboratory**
 - ☒ CMP
 - ☒ TSH
- Medications**
 - ☒ Med PLACEHOLDER 1
 - ☒ JEMPERLI 500 mg injection, for intravenous use.
Dose 1-4: 500 mg every 3 weeks. Subsequent dosing beginning 3 weeks after Dose 4
Dose 5 onwards: 1,000 mg every 6 weeks
- Patient Education**
 - ☒ About Endometrial Cancer

Hypothetical Order Set Shown in the PowerPlan

INDICATIONS

- JEMPERLI, in combination with carboplatin and paclitaxel, followed by JEMPERLI as a single agent, is indicated for the treatment of adult patients with primary advanced or recurrent endometrial cancer (EC).
- JEMPERLI, as a single agent, is indicated for the treatment of adult patients with mismatch repair deficient (dMMR) recurrent or advanced:
 - EC, as determined by an FDA-approved test, that has progressed on or following prior treatment with a platinum-containing regimen in any setting and are not candidates for curative surgery or radiation, or
 - solid tumors, as determined by an FDA-approved test, that have progressed on or following prior treatment and who have no satisfactory alternative treatment options. This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

IMPORTANT SAFETY INFORMATION

Severe and Fatal Immune-Mediated Adverse Reactions

- Immune-mediated adverse reactions, which can be severe or fatal, can occur in any organ system or tissue and can occur at any time during or after treatment with a PD-1/PD-L1 –blocking antibody, including JEMPERLI.
- Monitor closely for signs and symptoms of immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function tests at baseline and periodically during treatment. For suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.

Severe and Fatal Immune-Mediated Adverse Reactions (cont.)

- Based on the severity of the adverse reaction, withhold or permanently discontinue JEMPERLI. In general, if JEMPERLI requires interruption or discontinuation, administer systemic corticosteroids (1 to 2 mg/kg/day prednisone or equivalent) until improvement to ≤Grade 1. Upon improvement to ≤Grade 1, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose immune-mediated adverse reaction is not controlled with corticosteroids.

Immune-Mediated Pneumonitis

- JEMPERLI can cause immune-mediated pneumonitis, which can be fatal. In patients treated with other PD-1/PD-L1 –blocking antibodies, the incidence of pneumonitis is higher in patients who have received prior thoracic radiation. Pneumonitis occurred in 2.3% (14/605) of patients, including Grade 2 (1.3%), Grade 3 (0.8%), and Grade 4 (0.2%) pneumonitis.

Immune-Mediated Colitis

- Colitis occurred in 1.3% (8/605) of patients, including Grade 2 (0.7%) and Grade 3 (0.7%) adverse reactions. Cytomegalovirus infection/reactivation have occurred in patients with corticosteroid-refractory immune-mediated colitis. In such cases, consider repeating infectious workup to exclude alternative etiologies.

Immune-Mediated Hepatitis

- JEMPERLI can cause immune-mediated hepatitis, which can be fatal. Grade 3 hepatitis occurred in 0.5% (3/605) of patients.

Immune-Mediated Endocrinopathies

- Adrenal Insufficiency
 - Adrenal insufficiency occurred in 1.2% (7/605) of patients, including Grade 2 (0.5%) and Grade 3 (0.7%). For Grade 2 or higher adrenal insufficiency, initiate symptomatic treatment per institutional guidelines, including hormone replacement as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.

IMPORTANT SAFETY INFORMATION (cont.)

Immune-Mediated Endocrinopathies (cont.)

- Hypophysitis
 - JEMPERLI can cause immune-mediated hypophysitis. Grade 3 hypophysitis occurred in 0.4% (1/241) of patients receiving JEMPERLI in combination with carboplatin and paclitaxel. Grade 2 hypophysitis occurred in 0.2% (1/605) of patients receiving JEMPERLI as a single agent. Initiate hormone replacement as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.
- Thyroid Disorders
 - Grade 2 thyroiditis occurred in 0.5% (3/605) of patients. Grade 2 hypothyroidism occurred in 12% (30/241) of patients receiving JEMPERLI in combination with carboplatin and paclitaxel. Grade 2 hypothyroidism occurred in 8% (46/605) of patients receiving JEMPERLI as a single agent. Hyperthyroidism occurred in 3.3% (8/241) of patients receiving JEMPERLI in combination with carboplatin and paclitaxel, including Grade 2 (2.9%) and Grade 3 (0.4%). Hyperthyroidism occurred in 2.3% (14/605) of patients receiving JEMPERLI as a single agent, including Grade 2 (2.1%) and Grade 3 (0.2%). Initiate thyroid hormone replacement or medical management of hyperthyroidism as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.
- Type 1 Diabetes Mellitus, Which Can Present with Diabetic Ketoacidosis
 - JEMPERLI can cause type 1 diabetes mellitus, which can present with diabetic ketoacidosis. Grade 3 type 1 diabetes mellitus occurred in 0.4% (1/241) of patients receiving JEMPERLI in combination with carboplatin and paclitaxel. Grade 3 type 1 diabetes mellitus occurred in 0.2% (1/605) of patients receiving JEMPERLI as a single agent. Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Initiate treatment with insulin as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.

Immune-Mediated Nephritis with Renal Dysfunction

- JEMPERLI can cause immune-mediated nephritis, which can be fatal. Grade 2 nephritis, including tubulointerstitial nephritis, occurred in 0.5% (3/605) of patients.

Immune-Mediated Dermatologic Adverse Reactions

- JEMPERLI can cause immune-mediated rash or dermatitis. Bullous and exfoliative dermatitis, including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug rash with eosinophilia and systemic symptoms (DRESS), have occurred with PD-1/PD-L1–blocking antibodies. Topical emollients and/or topical corticosteroids may be adequate to treat mild to moderate non-bullous/exfoliative rashes. Withhold or permanently discontinue JEMPERLI depending on severity.

Other Immune-Mediated Adverse Reactions

- The following clinically significant immune-mediated adverse reactions occurred in <1% of the 605 patients treated with JEMPERLI or were reported with the use of other PD-1/PD-L1–blocking antibodies. Severe or fatal cases have been reported for some of these adverse reactions.
 - *Nervous System:* Meningitis, encephalitis, myelitis and demyelination, myasthenic syndrome/myasthenia gravis, Guillain-Barré syndrome, nerve paresis, autoimmune neuropathy
 - *Cardiac/Vascular:* Myocarditis, pericarditis, vasculitis
 - *Ocular:* Uveitis, iritis, other ocular inflammatory toxicities. Some cases can be associated with retinal detachment. Various grades of visual impairment to include blindness can occur
 - *Gastrointestinal:* Pancreatitis, including increases in serum amylase and lipase levels, gastritis, duodenitis
 - *Musculoskeletal and Connective Tissue:* Myositis/polymyositis, rhabdomyolysis and associated sequelae including renal failure, arthritis, polymyalgia rheumatica

IMPORTANT SAFETY INFORMATION (cont.)

Other Immune-Mediated Adverse Reactions (cont.)

- *Endocrine*: Hypoparathyroidism
- *Other (Hematologic/Immune)*: Autoimmune hemolytic anemia, aplastic anemia, hemophagocytic lymphohistiocytosis, systemic inflammatory response syndrome, histiocytic necrotizing lymphadenitis (Kikuchi lymphadenitis), sarcoidosis, immune thrombocytopenia, solid organ transplant rejection, other transplant (including corneal graft) rejection

Infusion-Related Reactions

- Severe or life-threatening infusion-related reactions have been reported with PD-1/PD-L1 –blocking antibodies. Severe infusion-related reactions (Grade 3) occurred in 0.2% (1/605) of patients receiving JEMPERLI. Monitor patients for signs and symptoms of infusion-related reactions. Interrupt or slow the rate of infusion or permanently discontinue JEMPERLI based on severity of reaction.

Complications of Allogeneic HSCT

- Fatal and other serious complications can occur in patients who receive allogeneic hematopoietic stem cell transplantation (HSCT) before or after treatment with a PD-1/PD-L1 –blocking antibody, which may occur despite intervening therapy. Monitor patients closely for transplant-related complications and intervene promptly.

Embryo-Fetal Toxicity and Lactation

- Based on its mechanism of action, JEMPERLI can cause fetal harm. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with JEMPERLI and for 4 months after their last dose. Because of the potential for serious adverse reactions from JEMPERLI in a breastfed child, advise women not to breastfeed during treatment with JEMPERLI and for 4 months after their last dose.

Common Adverse Reactions

The most common adverse reactions ($\geq 20\%$), including laboratory abnormalities, in patients with EC who received JEMPERLI in combination with carboplatin and paclitaxel were decreased hemoglobin, increased creatinine, peripheral neuropathy, decreased white blood cell count, fatigue, nausea, alopecia, decreased platelets, increased glucose, decreased lymphocytes, decreased magnesium, decreased neutrophils, increased AST, arthralgia, rash, constipation, diarrhea, increased ALT, decreased potassium, decreased albumin, decreased sodium, increased alkaline phosphatase, abdominal pain, dyspnea, decreased appetite, increased amylase, decreased phosphate, urinary tract infection, and vomiting.

The most common adverse reactions ($\geq 20\%$) in patients with dMMR EC who received JEMPERLI as a single agent were fatigue/asthenia, anemia, nausea, diarrhea, constipation, vomiting, and rash. The most common Grade 3 or 4 laboratory abnormalities ($> 2\%$) were decreased lymphocytes, decreased sodium, increased alanine aminotransferase, increased creatinine, decreased neutrophils, decreased albumin, and increased alkaline phosphatase.

The most common adverse reactions ($\geq 20\%$) in patients with dMMR solid tumors who received JEMPERLI as a single agent were fatigue/asthenia, anemia, diarrhea, and nausea. The most common Grade 3 or 4 laboratory abnormalities ($\geq 2\%$) were decreased lymphocytes, decreased sodium, increased alkaline phosphatase, and decreased albumin.

Please see full [Prescribing Information](#), including [Medication Guide](#).

Trademarks are property of their respective owners.

GSK ©2024 GSK or licensor.
PMUS-DSTBROC240012 July 2024
Produced in USA.

File name: EMR Guide Oracle Health