



ADDING JEMPERLI (DOSTARLIMAB-GXLY) TO AN iKnowMed™ TREATMENT REGIMEN

For Your Appropriate Adult Patients
With dMMR Recurrent or Advanced:
Endometrial Cancer or Solid Tumors

INDICATIONS

JEMPERLI is indicated for the treatment of adult patients with mismatch repair deficient (dMMR) recurrent or advanced:

- endometrial cancer (EC), as determined by an FDA-approved test, that has progressed on or following prior treatment with a platinum-containing regimen in any setting and are not candidates for curative surgery or radiation, or
- solid tumors, as determined by an FDA-approved test, that have progressed on or following prior treatment and who have no satisfactory alternative treatment options. This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

IMPORTANT SAFETY INFORMATION

Severe and Fatal Immune-Mediated Adverse Reactions

- Immune-mediated adverse reactions, which can be severe or fatal, can occur in any organ system or tissue and can occur at any time during or after treatment with a PD-1/PD-L1 –blocking antibody, including JEMPERLI.
- Monitor closely for signs and symptoms of immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function tests at baseline and periodically during treatment. For suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.

Please see additional [Important Safety Information](#)
on pages 8-10 and full [Prescribing Information](#).

dMMR = mismatch repair deficient.
iKnowMed is a trademark of US Oncology, Inc.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

ABOUT THIS GUIDE

This electronic health record (EHR) guide is intended to help health care providers (HCP) who want to create treatment Regimens that include JEMPERLI (dostarlimab-gxly) or want to add JEMPERLI to an existing treatment Regimen. Treatment Regimens group together order sets for medications, lab testing, procedures, and other aspects of care based on the patient's diagnosis and condition. It is important to evaluate oncology treatment plans frequently as treatment options, such as JEMPERLI, become available.

This guide does not constitute guidance for treatment or medical advice. It is the responsibility of the HCP to select a treatment based on their independent medical judgment and the needs of each individual patient.

The examples and instructions listed in this guide are based on the most recent version of iKnowMed. Locations, illustrations, and terminology are subject to change with system updates. This guide is meant to serve as an overview only and should not replace detailed instructions provided to you by your internal or external EHR support resources. GSK makes no claims or warranties about the applicability or appropriateness of this information. This guide has not been reviewed or endorsed by iKnowMed. GSK does not endorse or recommend any EHR system.

EHR REGIMENS HELP SIMPLIFY ONCOLOGY WORKFLOWS

Treatment Regimens are commonly used to help facilitate patient care. After the initial release of a treatment Regimen by a medical society, the iKnowMed system may benefit from a clinical update. The optimization of treatment Regimens is a common process and provides an opportunity to incorporate treatment updates. Treatment Regimens are typically modified at the health system level to help reduce treatment variation. Typically, an oncology practice will conduct a clinical review process to confirm and approve a suggested Regimen optimization. Various stakeholders may participate in reviewing treatment Regimen optimization requests prior to the approval.

As new tests, treatments, and protocols evolve, it may be appropriate to adjust patients' existing treatment plans and monitoring. With appropriate permissions, users can modify existing treatment Regimens for ease in adding orders to an existing treatment Regimen.

As treatment options such as JEMPERLI become available, it may be necessary to create a new Regimen or to update an existing Regimen to remove system obstacles to prescribe JEMPERLI for its approved indications. Updating relevant Regimens to include JEMPERLI communicates to the care team that it is available to order for appropriate patients.

NOTE: If JEMPERLI is not available for selection in iKnowMed, the practice may need to run a drug database update. As a backup option, the practice EHR Support/IT Team may be able to manually add JEMPERLI, subject to the practice's business rules for drug database maintenance.

CREATING OR EDITING A TREATMENT REGIMEN

Upon request and approval from the Clinical Team, the practice IT Team creates treatment Regimens that include the necessary orders for a given course of treatment. When an HCP assigns a protocol to a patient, it becomes the patient's treatment Regimen. Treatment Regimens can be saved in an HCP's individual library—and can be optionally shared publicly with others in the practice or in the entire health system.

EDITING AN EXISTING REGIMEN

Creating a Regimen Template

Custom Regimen Templates can be created only with full permissions. However, a Regimen Template can be created using a copy of an existing Regimen. Copying a Regimen Template is the method described in this guide.

1. From the **Manage** menu, select **Regimen Templates**.
2. In the **Reference Name or Display Name** field, select the Regimen to copy from. For example, dMMR.

Admin's Dashboard		Regimen Templates ✕	
<div> <div>ADD REGIMEN</div> <div>EDIT</div> <div>AUDIT HISTORY</div> <div>REMOVE</div> </div>			
Reference Name or Display Name	Owner Practice	Last Modified By	Status
dMMR			ACTIVE
Problem Groups	Stages	Region	

Regimen Template Search

3. Select **Copy Regimen**.
4. Update the **Reference Name** (internal) and the **Display Name**; select **Save**.
5. From the **Regimen Search Rules & Problem Associations** section, search for and select the problems for which the Regimen is appropriate.
6. Define Regimen Rule detail: **Problem Groups**, **Region**, and **Location** as appropriate for the practice. Select **Save**.
7. Select **Add Rule**.

Admin's Dashboard		Regimen Templates		Regimen Name Here ✕	
<div> <div>SAVE</div> <div>SAVE AND INCREMENT VERSION</div> <div>COPY REGIMEN</div> <div>RESET</div> <div>PRINT</div> </div>					
Regimen Details					

Copy Regimen Option

Regimen Search Rules & Problem Associations					
Note: Changes in this section are automatically saved.					
Simple Regimen Rules		Advanced Regimen Rule			
Search:				ADD RULE	
Problem Groups	Region	Location	Regimen Names	Remove	

Assigning Problem Associations

Manage	Admin
Clinical Note Templates	
Clinical Trials	
Dispensable	
Inferences	
Labs/Imaging & Services	
Medications	
Order Sets	
Orderable Groups	
Problems	
Problem Groups	
Regimen Templates	
Supplies	

Example of the Manage, Regimen Menu Templates Option

Creating a New Regimen Template (cont.)

8. Complete Regimen details:
 - Select **Status** "Suspended" until testing is completed, then set to **Active**
 - Select the appropriate **Regimen Type**
9. Review and update patient education information URLs and reference citation links as appropriate and save the edits.
10. Select **Save** to save Regimen information before editing the medication information.

Entering Regimen Details

Updating the New Regimen

1. Remove items from the originally copied Regimen that are not appropriate for this Regimen.
2. In the **Regimen Items** list, add items to the Regimen. From the appropriate Treatment Group, select **+** on the right side of the Treatment Group.

C / P	Pre-Check	Name	
		IMMUNOTHERAPY	+
			+

Updating Regimen Items

Updating the New Regimen (cont.)

3. Search for and select JEMPERLI; select **Save** to add to the template.
4. Select the medication item and add medication details as appropriate.
5. Select **Formula Dose** to enter appropriate dosing instructions.
 - Administration: For Preparation and Administration, please refer to the [Prescribing Information](#)
6. Select **Save** to create the new Regimen.

C / P	Pre-Check	Name	
COPY PASTE			+ ADD GROUP
<input type="checkbox"/>	R	O	IMMUNOTHERAPY + x
<input type="checkbox"/>	<input checked="" type="checkbox"/>		JEMPERLI x
<input type="checkbox"/>			PREMEDICATIONS + x

Example of a Medication Added to the Regimen

JEMPERLI, IV
required

☐ Rx
☒ Formula Dose
 QUICK SIG PICK SHOW DRUG FORMS

Dose Unit
 ADJUST DOSE

JEMPERLI

Dose Unit Form
 500 mg Injection

Route Method Frequency
 Intravenous Select once ☐ PRN

Administer Over

Admix Fluid Volume
 Select mL

Instructions ☐ Instructions replace required fields
 Duration

Administer infusion intravenously over 30 minutes

- JEMPERLI injection, for intravenous use
- Dose 1 through 4: 500 mg every 3 weeks
- Subsequent dosing beginning 3 weeks after Dose 4 (Dose 5 onwards): 1,000 mg every 6 weeks
- Administration: For Preparation and Administration, please refer to the Prescribing Information

☐ Allow substitutions.
 ☐ Do Not Bill Drug
 ☐ Do Not Bill Administration

SAVE CANCEL

Adding Dosage Details and Instructions to the Regimen

Chart Summary	Clinical Profile	Flowsheet	Orders	Results	Documents	Demographics	Nursing Care
New Orders (0)	Medications (0)	Regimens (0)	Order Review	Order History	Drafts		
IMMUNOTHERAPY		Add an Order	Search Orderables	Dose			
JEMPERLI		JEMPERLI 500 mg/10 mL		500 mg/10 mL			
<input checked="" type="checkbox"/> 500 mg/10 mL intravenous, over 30 minutes, ONCE, then every 3 weeks for 4 cycles		<input type="checkbox"/> 1,000 mg, over 30 minutes, ONCE, every 6 weeks					
BIOSIMILAR		Add an Order	Search Orderables	Dose			

Example of the Regimen in the Patient Chart

Creating an Order Panel to Add to a New or Existing Regimen

1. Select **Manage, Order Sets** to create or add lab orders and patient education orders for dMMR.
2. Choose **Add Order Set**. Name the Order Set.
3. Define the panel, using detail below, then **Save**.
 - Type of **Order Panel**
 - Order type of **New Orders**

Example of Creating an Order Panel

4. Select the newly created **Order Panel** to add appropriate items. Select **Add Items**.
5. Search for and select lab orders.
6. Include other orders as desired.
7. Select **Save**.

Example of Orderables Search

INDICATIONS

JEMPERLI is indicated for the treatment of adult patients with mismatch repair deficient (dMMR) recurrent or advanced:

- endometrial cancer (EC), as determined by an FDA-approved test, that has progressed on or following prior treatment with a platinum-containing regimen in any setting and are not candidates for curative surgery or radiation, or
- solid tumors, as determined by an FDA-approved test, that have progressed on or following prior treatment and who have no satisfactory alternative treatment options. This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

IMPORTANT SAFETY INFORMATION

Severe and Fatal Immune-Mediated Adverse Reactions

- Immune-mediated adverse reactions, which can be severe or fatal, can occur in any organ system or tissue and can occur at any time during or after treatment with a PD-1/PD-L1–blocking antibody, including JEMPERLI.
- Monitor closely for signs and symptoms of immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function tests at baseline and periodically during treatment. For suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.
- Based on the severity of the adverse reaction, withhold or permanently discontinue JEMPERLI. In general, if JEMPERLI requires interruption or discontinuation, administer systemic corticosteroids (1 to 2 mg/kg/day prednisone or equivalent) until improvement to \leq Grade 1. Upon improvement to \leq Grade 1, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose immune-mediated adverse reaction is not controlled with corticosteroids.

Immune-Mediated Pneumonitis

- JEMPERLI can cause immune-mediated pneumonitis, which can be fatal. In patients treated with other PD-1/PD-L1–blocking antibodies, the incidence of pneumonitis is higher in patients who have received prior thoracic radiation. Pneumonitis occurred in 2.3% (14/605) of patients, including Grade 2 (1.3%), Grade 3 (0.8%), and Grade 4 (0.2%) pneumonitis.

Immune-Mediated Colitis

- Colitis occurred in 1.3% (8/605) of patients, including Grade 2 (0.7%) and Grade 3 (0.7%) adverse reactions. Cytomegalovirus infection/reactivation have occurred in patients with corticosteroid-refractory immune-mediated colitis. In such cases, consider repeating infectious workup to exclude alternative etiologies.

Immune-Mediated Hepatitis

- JEMPERLI can cause immune-mediated hepatitis, which can be fatal. Grade 3 hepatitis occurred in 0.5% (3/605) of patients.

Immune-Mediated Endocrinopathies

- Adrenal Insufficiency
 - Adrenal insufficiency occurred in 1.2% (7/605) of patients, including Grade 2 (0.5%) and Grade 3 (0.7%). For Grade 2 or higher adrenal insufficiency, initiate symptomatic treatment per institutional guidelines, including hormone replacement as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.
- Hypophysitis
 - JEMPERLI can cause immune-mediated hypophysitis. Grade 2 hypophysitis occurred in 0.2% (1/605) of patients. Initiate hormone replacement as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.
- Thyroid Disorders
 - Grade 2 thyroiditis occurred in 0.5% (3/605) of patients. Grade 2 hypothyroidism occurred in 7.6% (46/605) of patients. Hyperthyroidism occurred in 2.3% (14/605) of patients, including Grade 2 (2.1%) and Grade 3 (0.2%). Initiate hormone replacement or medical management of hyperthyroidism as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.

IMPORTANT SAFETY INFORMATION (cont.)

Immune-Mediated Endocrinopathies (cont.)

- Type 1 Diabetes Mellitus, Which Can Present with Diabetic Ketoacidosis
 - JEMPERLI can cause type 1 diabetes mellitus, which can present with diabetic ketoacidosis. Grade 3 type 1 diabetes mellitus occurred in 0.2% (1/605) of patients. Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Initiate treatment with insulin as clinically indicated. Withhold or permanently discontinue JEMPERLI depending on severity.

Immune-Mediated Nephritis with Renal Dysfunction

- JEMPERLI can cause immune-mediated nephritis, which can be fatal. Grade 2 nephritis, including tubulointerstitial nephritis, occurred in 0.5% (3/605) of patients.

Immune-Mediated Dermatologic Adverse Reactions

- JEMPERLI can cause immune-mediated rash or dermatitis. Bullous and exfoliative dermatitis, including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug rash with eosinophilia and systemic symptoms (DRESS), have occurred with PD-1/PD-L1–blocking antibodies. Topical emollients and/or topical corticosteroids may be adequate to treat mild to moderate non-bullous/exfoliative rashes. Withhold or permanently discontinue JEMPERLI depending on severity.

Other Immune-Mediated Adverse Reactions

- The following clinically significant immune-mediated adverse reactions occurred in <1% of the 605 patients treated with JEMPERLI or were reported with the use of other PD-1/PD-L1–blocking antibodies. Severe or fatal cases have been reported for some of these adverse reactions.
 - *Nervous System:* Meningitis, encephalitis, myelitis and demyelination, myasthenic syndrome/myasthenia gravis, Guillain-Barre syndrome, nerve paresis, autoimmune neuropathy
 - *Cardiac/Vascular:* Myocarditis, pericarditis, vasculitis
 - *Ocular:* Uveitis, iritis, other ocular inflammatory toxicities. Some cases can be associated with retinal detachment. Various grades of visual

impairment to include blindness can occur

- *Gastrointestinal:* Pancreatitis, including increases in serum amylase and lipase levels, gastritis, duodenitis
- *Musculoskeletal and Connective Tissue:* Myositis/polymyositis, rhabdomyolysis and associated sequelae including renal failure, arthritis, polymyalgia rheumatica
- *Endocrine:* Hypoparathyroidism
- *Other (Hematologic/Immune):* Autoimmune hemolytic anemia, aplastic anemia, hemophagocytic lymphohistiocytosis, systemic inflammatory response syndrome, histiocytic necrotizing lymphadenitis (Kikuchi lymphadenitis), sarcoidosis, immune thrombocytopenia, solid organ transplant rejection

Infusion-Related Reactions

- Severe or life-threatening infusion-related reactions have been reported with PD-1/PD-L1–blocking antibodies. Severe infusion-related reactions (Grade 3) occurred in 0.2% (1/605) of patients receiving JEMPERLI. Monitor patients for signs and symptoms of infusion-related reactions. Interrupt or slow the rate of infusion or permanently discontinue JEMPERLI based on severity of reaction.

Complications of Allogeneic HSCT

- Fatal and other serious complications can occur in patients who receive allogeneic hematopoietic stem cell transplantation (HSCT) before or after treatment with a PD-1/PD-L1–blocking antibody, which may occur despite intervening therapy. Monitor patients closely for transplant-related complications and intervene promptly.

Embryo-Fetal Toxicity and Lactation

- Based on its mechanism of action, JEMPERLI can cause fetal harm. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with JEMPERLI and for 4 months after their last dose. Because of the potential for serious adverse reactions from JEMPERLI in a breastfed child, advise women not to breastfeed during treatment with JEMPERLI and for 4 months after their last dose.

IMPORTANT SAFETY INFORMATION (cont.)

Common Adverse Reactions

The most common adverse reactions ($\geq 20\%$) in patients with dMMR EC were fatigue/asthenia, anemia, rash, nausea, diarrhea, constipation, and vomiting. The most common Grade 3 or 4 laboratory abnormalities ($> 2\%$) were decreased lymphocytes, decreased sodium, increased alanine aminotransferase, increased creatinine, decreased neutrophils, decreased albumin, and increased alkaline phosphatase.

The most common adverse reactions ($\geq 20\%$) in patients with dMMR solid tumors were fatigue/asthenia, anemia, diarrhea, and nausea. The most common Grade 3 or 4 laboratory abnormalities ($\geq 2\%$) were decreased lymphocytes, decreased sodium, increased alkaline phosphatase, and decreased albumin.

Please see full [Prescribing Information](#).

Trademarks are property of their respective owners.



©2023 GSK or licensor.
DSTOGM230023 March 2023
Produced in USA. 0002-0022-94

File name: EHR Guide iKnowMed